Fractal arithmetic and calculus on Sierpiński sets

Diederik Aerts[1], Marek Czachor[1,2], and Maciej Kuna[1,2]
[1] *Centrum Leo Apostel (CLEA), Vrije Universiteit Brussel, 1050 Brussels, Belgium,*
[2] *Wydział Fizyki Technicznej i Matematyki Stosowanej, Politechnika Gdańska, 80-233 Gdańsk, Poland*

Calculus that starts with elementary non-Diophantine arithmetic operations of a Burgin type can be naturally incorporated into fractals of a Sierpiński type. Non-Diophantine definitions of derivatives and integrals are much simpler from what one finds in the more traditional literature of the subject, and automatically lead to harmonic analysis of functions defined on fractals. As an application we perform a Fourier transform of a real-valued function with Sierpiński-set domain.

PACS numbers: 04.60.-m, 05.45.Df, 02.20.Qs

arXiv:1606.01337v3 [math.GN] 20 Mar 2017

I. INTRODUCTION

Apparently, the first attempt of a Fourier-type analysis on fractals can be found in studies of diffusion on Sierpiński gaskets [1, 2]. A generator of the diffusion process plays there the same role as a Laplacian on a manifold, so the corresponding eigenfunction expansion may be regarded as a form of harmonic analysis. An alternative route to eigenfunction expansions on fractals is to define Laplacians or gradients more directly. Here certain approaches begin with Dirichlet forms on self-similar fractals, or one takes as a departure point discrete Laplacians and performes an appropriate limit [3–6]. Four alternative definitions of a gradient (due to Kusuoka, Kigami, Strichartz and Teplyaev) are discussed in this context in [7]. Self-similarity is typically an important technical assumption. Although Laplacians defined in the above ways cannot be regarded as second-order operators, an approach where Laplacians are indeed second-order is nevertheless possible and was introduced by Fujita [8, 9] and further developed by a number of authors [10–14].

A second traditional approach to harmonic analysis on fractals comes from the notion of self-similar fractal measures. The classic result of Jorgensen and Pedersen [15] states that the method works for certain fractals, such as the quaternary Cantor set, but fails in the important case of the ternary middle-third Cantor set. Moreover, the method is inapplicable in realistic cases of non-self-similar fractals.

Quite recently we have shown [16] that calculus based on Burgin's non-Diophantine arithmetic [17–20] leads to a simple and very efficient construction of a Fourier transform on fractals of a Cantor type. Gradients and Laplacians are here, respectively, first- and second-order differential operators, and self-similarity plays no role whatsoever. There is completely no difficulty with Fourier analysis of functions mapping arbitrary Cantor sets into themselves, so Jorgensen-Pedersen-type restrictions are no longer valid.

The goal of the present paper is to show that the non-Diophantine formalism applies also to fractals of a Sierpiński type, an example of great importance for fractal analysis. We explicitly show how to perform the construction on a double cover of a Sierpiński set. Similarly to the Cantor case, self-similarity is inessential for the formalism. What is essential, however, is the existence of a bijection f between the fractal in question and $\mathbb{R}$.

In the Sierpiński case the bijection has properties reminiscent of Peano curves [21]. The very idea that there are links between Sierpiński-type fractals and space-filling curves is not new, and was used by Molitor *et al.* in [22] in their construction of Laplacians on fractals. However, in all other respects the approach from [22] is different from what we discuss below. The idea of employing a one-dimensional integration for finding higher dimensional integrals is known [23], but apparently has not been used in fractal contexts so far.

Since any fractal whose cardinality is continuum can be equipped with a bijection mapping it into $\mathbb{R}$, the construction is quite universal. From a practical perspective, the only difficulty is to find the bijection explicitly, but once we have found it the remaining procedure is systematic and easy to work with.

II. SIERPIŃSKI SET

Consider $x \in \mathbb{R}_+$ and its ternary representation $x = (t_n \dots t_0.t_{-1}t_{-2}\dots)_3$. If x has two different ternary representations, we choose the one that ends with infinitely many 2s. Keeping the digits unchanged let us change the base from 3 to 4, i.e.

$$x = (t_n \dots t_0.t_{-1}t_{-2}\dots)_3 \mapsto (t_n \dots t_0.t_{-1}t_{-2}\dots)_4 = y. \quad (1)$$

The quaternary representation of y is unique, and it does not involve the digit 3. Next, let us parametrize the quaternary digits in a binary way, but written in a column form: $0 = \begin{smallmatrix}0\\0\end{smallmatrix}$, $1 = \begin{smallmatrix}0\\1\end{smallmatrix}$, $2 = \begin{smallmatrix}1\\0\end{smallmatrix}$, $3 = \begin{smallmatrix}1\\1\end{smallmatrix}$. y has been converted into a pair of binary sequences,

$$(t_n \dots t_0.t_{-1}t_{-2}\dots)_4 \mapsto \begin{pmatrix} a_n \dots a_0.a_{-1}a_{-2}\dots \\ b_n \dots b_0.b_{-1}b_{-2}\dots \end{pmatrix}_2 \quad (2)$$

where $(a_j, b_j) \neq (1, 1)$ for any j. The resulting sequences are in a one-to-one relation with the x we have started with. Each of the two sequences defines a number in

binary notation: we have mapped x into a point of the plane $\mathbb{R}_+ \times \mathbb{R}_+$. The image of $\mathbb{R}_+$ under our algorithm defines a Sierpiński-type set. The algorithm is not invertible. Indeed, take the point $(1,1)$. Depending on the way we represent it binarily we find

$$\begin{pmatrix} 1.(0) \\ 0.(1) \end{pmatrix}_2 \mapsto \big(2.(1)\big)_4 \mapsto \big(2.(1)\big)_3 = 2.5,$$

and

$$\begin{pmatrix} 0.(1) \\ 1.(0) \end{pmatrix}_2 \mapsto \big(1.(2)\big)_4 \mapsto \big(1.(2)\big)_3 = 2.$$

The ambiguity comes from the two identifications

$$(1,1) = \begin{pmatrix} 1.(0) \\ 0.(1) \end{pmatrix}_2, \quad (1,1) = \begin{pmatrix} 0.(1) \\ 1.(0) \end{pmatrix}_2. \tag{3}$$

However, if we write the above two relations as

$$(1,1)_- = \begin{pmatrix} 1.(0) \\ 0.(1) \end{pmatrix}_2, \tag{4}$$

$$(1,1)_+ = \begin{pmatrix} 0.(1) \\ 1.(0) \end{pmatrix}_2, \tag{5}$$

and treat the two points $(1,1)_\pm$ as belonging to two different sides of an oriented plane, the ambiguity of the inverse alorithm disappears. The relation

$$(1,1)_- \leftrightarrow 2.5, \quad (1,1)_+ \leftrightarrow 2 \tag{6}$$

is one-to-one.

Let us therefore index with '+' (resp. '−') those pairs (a,b) where a is a rational number represented by a binary sequence involving $(1)_2$ (resp. $(0)_2$), and b is a rational number whose binary representation contains $(0)_2$ (resp. $(1)_2$). In both cases $(a_j, b_j) \neq (1,1)$ by construction. The corresponding rational numbers involve, respectively, $(2)_3$ and $(1)_3$.

But what about the other cases, such as a, b irrational, or a irrational but b rational? It turns out that the ambiguity is absent. In order to prove it, first of all note that we did not have to consider the cases

$$(1,1) = \begin{pmatrix} 0.(1) \\ 0.(1) \end{pmatrix}_2, \quad (1,1) = \begin{pmatrix} 1.(0) \\ 1.(0) \end{pmatrix}_2, \tag{7}$$

since the pairs $(a_j, b_j) = (1,1)$ cannot appear as a result of the algorithm, and two infinite sequences of 0s would imply that $(t_n \dots t_0.t_{-1}t_{-2}\dots)_4$ ends with an infinite sequence 0s, a form excluded by the algorithm.

The same mechanism eliminates all the remaining ambiguities:

(A) If a, b are both irrational, or a is irrational and b rational-periodic, their binary forms are unique.

(B) If a is irrational (or rational-periodic), but b rational non-periodic, then b cannot end with infinitely many 1s, as it would mean that a ends with infinitely many 0s. So these cases are again unique. Conclusions are unchanged if one interchanges a and b.

(C) The only ambiguity appears if a ends with infinitely many 0s, but b with infinitely many 1s (or the other way around). But this is the case we have started with.

In cases (A) and (B) we identify $(a,b)_+ = (a,b)_- = (a,b)$. Only the (countable) case (C) requires a two-sided plane $(a,b)_+ \neq (a,b)_-$. The case (C) occurs for those $x \in \mathbb{R}$ whose ternary representation ends with $(2)_3$ or $(1)_3$. Only the latter numbers are mapped into $(a,b)_-$.

As we can see, what we have constructed is a version of a double cover of the Sierpiński set.

Our algorithm defines an injective map g_+ of $\mathbb{R}_+$ into a two-sided plane, with the above mentioned identifications. Let us extend g_+ to g by $g(|x|) = g_+(|x|)$, $g(-|x|) = -g_+(|x|)$. The image $S = g(\mathbb{R})$ is our definition of the Sierpiński set. Denoting $f = g^{-1}$, $f : S \to \mathbb{R}$ we obtain Burgin's arithmetic intrinsic to S,

$$x \oplus y = f^{-1}\big(f(x) + f(y)\big), \tag{8}$$

$$x \ominus y = f^{-1}\big(f(x) - f(y)\big), \tag{9}$$

$$x \odot y = f^{-1}\big(f(x) f(y)\big), \tag{10}$$

$$x \oslash y = f^{-1}\big(f(x) / f(y)\big). \tag{11}$$



FIG. 1: The inverse image $f^{-1}([0,1))$, view from the positive side of the oriented plane.

Fig. 1 shows the set $f^{-1}\big([0,1)\big)$. The set is self-similar and its Hausdorff dimension is $\log_2 3$.

III. ARITHMETIC ON THE SIERPIŃSKI SET

Let us begin with neutral elements of addition and multiplication in S. By definition, $0' \oplus x = x$, $1' \odot x = x$,



FIG. 2: The image of the first 200 natural numbers, $f^{-1}(\{1,\ldots,200\})$. All natural numbers are mapped into the positive side of the oriented plane.

where

$$0' = f^{-1}(0) = (0,0) \in S, \quad (12)$$
$$1' = f^{-1}(1) = (1,0)_+ \in S. \quad (13)$$

In non-Diophantine arithmetic multiplication is a repeated addition in the following sense [16]. Let $n \in \mathbb{N}$ and $n' = f^{-1}(n) \in S$. Then

$$n' \oplus m' = (n+m)', \quad (14)$$
$$n' \odot m' = (nm)' \quad (15)$$
$$= \underbrace{m' \oplus \cdots \oplus m'}_{n\text{times}}. \quad (16)$$

A power function $A(x) = x \odot \cdots \odot x$ (n times) is denoted by $x^{n'}$, which is consistent with

$$x^{n'} \odot x^{m'} = x^{(n+m)'} = x^{n' \oplus m'}. \quad (17)$$

All Sierpińskian integers are represented by pairs of integers, a representation somewhat similar to complex numbers, but with different rules of addition and multiplication, as illustrated by

$$3' \oplus 4' = (2,0)_+ \oplus (1,2)_+ = 7' = (3,0)_+ \quad (18)$$

(numbers represented in decimal form).

IV. CALCULUS

The derivative of a function $A : S \to S$ is defined by [16, 17, 20]

$$\frac{DA(x)}{Dx} = \lim_{h \to 0'} \Big(A(x \oplus h) \ominus A(x) \Big) \oslash h. \quad (19)$$

A Laplacian on S is just the second derivative

$$\Delta A(x) = \frac{D}{Dx} \frac{DA(x)}{Dx}. \quad (20)$$

Our Δ differs from the other definitions of Laplacians on Sierpiński sets occurring in the literature [5, 6], but is simple and easy to work with. The formalism from [16, 17, 20] can be applied here with no modification, including integration, complex numbers, harmonic analysis, differential equations...

What was *not* explained in [16, 17, 20] was how to proceed with functions that do not map the fractal in question into itself. So, consider two sets, $\mathbb{X}$ and $\mathbb{Y}$ say, equipped with bijections $f_{\mathbb{Y}} : \mathbb{Y} \to \mathbb{R}$ and $f_{\mathbb{X}} : \mathbb{X} \to \mathbb{R}$, and arithmetics $\{\oplus_{\mathbb{Y}}, \odot_{\mathbb{Y}} : \mathbb{Y} \times \mathbb{Y} \to \mathbb{Y}\}$, $\{\oplus_{\mathbb{X}}, \odot_{\mathbb{X}} : \mathbb{X} \times \mathbb{X} \to \mathbb{X}\}$, defined by $f_{\mathbb{Y}}$ and $f_{\mathbb{X}}$. The bijection $f = f_{\mathbb{Y}}^{-1} \circ f_{\mathbb{X}} : \mathbb{X} \to \mathbb{Y}$ makes it possible to consider derivatives of functions $A : \mathbb{X} \to \mathbb{Y}$. Let $0'_{\mathbb{X}} = f_{\mathbb{X}}^{-1}(0)$ be the neutral element of addition in $\mathbb{X}$. We define

$$\frac{DA(x)}{Dx} = \lim_{h \to 0'_{\mathbb{X}}} \Big(A(x \oplus_{\mathbb{X}} h) \ominus_{\mathbb{Y}} A(x) \Big) \oslash_{\mathbb{Y}} f(h) \quad (21)$$
$$= f_{\mathbb{Y}}^{-1} \left(\frac{d}{df_{\mathbb{X}}(x)} \underbrace{f_{\mathbb{Y}} \circ A \circ f_{\mathbb{X}}^{-1}}_{a} \big[f_{\mathbb{X}}(x) \big] \right). \quad (22)$$

Details of the transition from (21) to (22) can be found in the Appendix. Accordingly,

$$\frac{DA(x)}{Dx} = f_{\mathbb{Y}}^{-1} \circ a' \circ f_{\mathbb{X}}(x) \quad (23)$$

where $a'(x) = \lim_{h \to 0} \big(a(x+h) - a(x) \big)/h$.

The integral is defined in a way guaranteeing the fundamental laws of calculus, relating derivatives and integrals. So, let $A : \mathbb{X} \to \mathbb{Y}$, and

$$\int_Y^X A(x) Dx = f_{\mathbb{Y}}^{-1} \left(\int_{f_{\mathbb{X}}(Y)}^{f_{\mathbb{X}}(X)} f_{\mathbb{Y}} \circ A \circ f_{\mathbb{X}}^{-1}(x) dx \right) \quad (24)$$

where $\int a(x)dx$ is the usual (say, Lebesgue) integral of a function $a : \mathbb{R} \to \mathbb{R}$.

In the Appendix we prove that

$$\frac{D}{DX} \int_Y^X A(x) Dx = A(X), \quad (25)$$
$$\int_Y^X \frac{DA(x)}{Dx} Dx = A(X) \ominus_{\mathbb{Y}} A(Y). \quad (26)$$

In the next section we apply the formalism to the problem of harmonic analysis on Sierpiński sets.

V. EXAMPLE: FOURIER TRANSFORM ON S

Let $\mathbb{X} = S$ and $\mathbb{Y} = \mathbb{R}$. Consider the function $A : S \to \mathbb{R}$ (Fig. 3),

$$A(x) = \begin{cases} 1 & \text{for } x \in S \cap (0,1)^2 \\ -1 & \text{for } x \in S \cap (-1,0)^2 \\ 0 & \text{otherwise} \end{cases} \quad (27)$$

Since $S \cap (-1,0]^2 = f_S^{-1}\big((-1,0]\big)$, $S \cap [0,1)^2 = f_S^{-1}\big([0,1)\big)$, $S \cap \{(0,0)\} = f_S^{-1}\big(0\big)$, we introduce $a : \mathbb{R} \to \mathbb{R}$,

$$a(x) = \begin{cases} 1 & \text{for } x \in (0,1) \\ -1 & \text{for } x \in (-1,0) \\ 0 & \text{otherwise} \end{cases} \tag{28}$$

Employing $f_{\mathbb{X}} = f_S$, $f_{\mathbb{Y}} = \mathrm{id}_{\mathbb{R}}$, we get

$$a = f_{\mathbb{Y}} \circ A \circ f_{\mathbb{X}}^{-1} = A \circ f_S^{-1}. \tag{29}$$

In order to perform Fourier analysis of A we have to introduce the basis of sines and cosines along the lines of [16], but adapted to the present context. The scalar product of two functions $G_j : S \to \mathbb{R}$, $G_j = g_j \circ f_S$, $g_j : \mathbb{R} \to \mathbb{R}$, $j = 1, 2$, reads

$$\langle G_1 | G_2 \rangle = \int_{\ominus_{\mathbb{Y}} T}^{T} G_1(x) \odot_{\mathbb{Y}} G_2(x) Dx \tag{30}$$

$$= f_{\mathbb{Y}}^{-1} \left(\int_{-f_S(T)}^{f_S(T)} g_1(x) g_2(x) dx \right) \tag{31}$$

$$= \int_{-1}^{1} g_1(x) g_2(x) dx = \langle g_1 | g_2 \rangle \tag{32}$$

where $\ominus_S T = 0'_S \ominus_S T = f_S^{-1}(-1)$. In our case $T = 1'_S = f_S^{-1}(1)$ (the neutral element of multiplication in S). The fact that $\langle g_1 | g_2 \rangle$ and $\langle G_1 | G_2 \rangle$ involve the same symbol of the scalar product will not lead to ambiguities.

Denoting

$$c_n(y) = \cos n\pi y, \quad n > 0 \tag{33}$$

$$s_n(y) = \sin n\pi y, \quad n > 0 \tag{34}$$

$$c_0(y) = 1/\sqrt{2}, \tag{35}$$

$$s_0(y) = 0, \tag{36}$$

$$C_n(x) = c_n\big(f_S(x)\big), \tag{37}$$

$$S_n(x) = s_n\big(f_S(x)\big), \tag{38}$$

we can apply the standard resolution of unity,

$$\delta(x - y) = \sum_{n \geq 0} \Big(c_n(x) c_n(y) + s_n(x) s_n(y) \Big), \tag{39}$$

and finally obtain

$$A(x) = \sum_{n \geq 0} \Big(C_n(x) \langle C_n | A \rangle + S_n(x) \langle S_n | A \rangle \Big) \tag{40}$$

$$= \sum_{n \geq 0} \Big(C_n(x) \langle c_n | a \rangle + S_n(x) \langle s_n | a \rangle \Big) \tag{41}$$

$$= \sum_{n > 0} \frac{2\,(1 - (-1)^n)}{n\pi} S_n(x) \tag{42}$$

Fig. 5 and Fig. 6 illustrate the reconstruction formula (42) with 5 and 50 Fourier terms, respectively. Figures 3 and 4 illustrate finite-sum Fourier reconstructions of the



FIG. 3: Function $A(x)$ defined in (27).

function plotted in Fig. 3. The negative-side of S occurs only as the image of those $x \in \mathbb{R}$ whose ternary representation ends with infinitely many 1s, whereas all finite-ternary-digit numbers are mapped into the positive side of S. This leads to the practical moral: Plotting functions with domains in S we can concentrate exclusively on the positive side of S, unless one employs a symbolic algorithm that recognizes numbers that end with $(1)_3$.

VI. CONCLUSION

The formalism proposed in [17] and further developed in [16, 20] naturally extends to a quite general class of fractals, including those of a Sierpiński type. The only nontrivial element of the construction is to find an explicit f. For a fractal whose cardinality equals continuum the existence of f follows directly from the fact that the cardinality of $\mathbb{R}$ is the same. Although one may, perhaps, have doubts as to the physical meaning of the derivative we employ, the associated integral is very practical from the point of view of harmonic analysis.

[1] S. Kusuoka, A diffusion process on a fractal, Probabilistic Methods in Mathematical Physics, Proceedings of Taniguchi International Symposium, K. Ito and N. Ikeda,



FIG. 4: Finite-sum reconstruction of $A(x)$ with 5 Fourier terms in (42) ...



FIG. 5: ...and with 50 terms. The Gibbs phenomenon is clearly visible.

eds., 251-274, Kinokuniya (1987).

[2] S. Goldstein, Random walks and diffusions on fractals, Percolation theory and ergodic theory of infinite particle systems, H. Kesten, ed., The IMA Volumes in Mathematics and its Applications, vol. 8, 121-129, Springer (1987).

[3] S. Kusuoka, Dirichlet forms on fractals and products of random matrices, Publ. Res. Inst. Math. Sci. **25**, 659-680 (1989).

[4] J. Kigami, A harmonic calculus on the Sierpinski spaces, Japan J. Appl. Math. **6**, 259-290 (1989).

[5] J. Kigami, Analysis on Fractals, Cambridge Tracts in Mathematics, vol. 143, Cambridge University Press, Cambridge (2001).

[6] R. S. Strichartz, Differential Equations on Fractals, Princeton University Press, Princeton (2006).

[7] A. Teplyaev, Gradients on fractals, J. Funct. Anal. **174**, 128-154 (2000).

[8] T. Fujita, A fractional dimension, self-similarity and a generalized diffusion operator, Probabilistic Methods in Mathematical Physics, Proceedings of Taniguchi International Symposium, K. Ito and N. Ikeda, eds., Kinokuniya, 1987, pp. 83-90.

[9] T. Fujita, Some asymptotic estimates of transition probability densities for generalized diffusion processes with self-similar speed measures, Publ. Res. Inst. Math. Sci. **26**, 819-840 (1990).

[10] U. Freiberg and M. Zähle, Harmonic calculus on fractals: A measure geometric approach (I), Potential Analysis **16**, 265-277 (2002).

[11] U. Freiberg, A survey on measure geometric Laplacians on Cantor like sets. Arabian J. Sci. Eng. **28**, 189-198 (2003).

[12] M. Zähle, Harmonic calculus on fractals: A measure geometric approach (II), Trans. Am. Math. Soc. **357**, 3407-3423 (2005).

[13] P. Arzt, Measure theoretic trigonometric functions, arXiv:1405.4693 [math.SP] (2014).

[14] M. Kesseböhmer, T. Samuel and H. Weyer, A note on measure-geometric Laplacians, arXiv:1411.2491 [math.FA] (2014).

[15] P. E. T. Jorgensen and S. Pedersen, Dense analytic subspaces in fractal L^2 spaces, J. Anal. Math. **75**, 185-228 (1998); preprint arXiv:funct-an/9709007.

[16] D. Aerts, M. Czachor, and M. Kuna, Fourier transforms on Cantor sets: A study in non-Diophantine arithmetic and calculus, Chaos, Solitons and Fractals **91**, 461468 (2016); arXiv:1603.05471 [math-ph] (2016).

[17] M. Czachor, Relativity of arithmetic as a fundamental symmetry of physics, Quantum Stud.: Math. Found. **3**, 123-133 (2016); arXiv:1412.8583 [math-ph] (2014).

[18] M. Burgin, Non-Diophantine Arithmetics, Ukrainian Academy of Information Sciences, Kiev (1997) (in Russian).

[19] M. Burgin, Introduction to projective arithmetics, arXiv:1010.3287 [math.GM] (2010).

[20] D. Aerts, M. Czachor, and M. Kuna, Crystallization of space: Space-time fractals from fractal arithmetic, Chaos, Solitons and Fractals **83**, 201-211 (2016).

[21] G. Peano, Sur une courbe, qui remplit toute une aire plane, Mathematische Annalen **36** (1): 15716 (1890).

[22] D. Molitor, N. Ott, and R. Strichartz, Using Peano curves to construct Laplacians on fractals, arXiv:1402.2106 [math.FA] (2014).

[23] N. Wiener, The Fourier Integral and Certain of its Appli-

cations, Cambridge University Press, Cambridge (1933).

Appendix

Here we give detailed proofs of formulas (21), (25), and (26).

$$\frac{DA(x)}{Dx} = \lim_{h\to 0'_{\mathbb{X}}} \Big(A(x\oplus_{\mathbb{X}} h)\ominus_{\mathbb{Y}} A(x)\Big)\oslash_{\mathbb{Y}} f(h) \tag{43}$$

$$= \lim_{h\to 0'_{\mathbb{X}}} f_{\mathbb{Y}}^{-1}\left(\frac{f_{\mathbb{Y}}\big(A(x\oplus_{\mathbb{X}} h)\ominus_{\mathbb{Y}} A(x)\big)}{f_{\mathbb{Y}}\big(f(h)\big)}\right) \tag{44}$$

$$= \lim_{h\to 0'_{\mathbb{X}}} f_{\mathbb{Y}}^{-1}\left(\frac{f_{\mathbb{Y}}\big(A(x\oplus_{\mathbb{X}} h)\ominus_{\mathbb{Y}} A(x)\big)}{f_{\mathbb{X}}(h)}\right) \tag{45}$$

$$= \lim_{h\to 0'_{\mathbb{X}}} f_{\mathbb{Y}}^{-1}\left(\frac{f_{\mathbb{Y}}\big(A(x\oplus_{\mathbb{X}} h)\big) - f_{\mathbb{Y}}\big(A(x)\big)}{f_{\mathbb{X}}(h)}\right) \tag{46}$$

$$= \lim_{h\to 0'_{\mathbb{X}}} f_{\mathbb{Y}}^{-1}\left(\frac{f_{\mathbb{Y}}\Big(A\Big(f_{\mathbb{X}}^{-1}\big[f_{\mathbb{X}}(x)+f_{\mathbb{X}}(h)\big]\Big)\Big) - f_{\mathbb{Y}}\Big(A\Big(f_{\mathbb{X}}^{-1}\big[f_{\mathbb{X}}(x)\big]\Big)\Big)}{f_{\mathbb{X}}(h)}\right) \tag{47}$$

$$= \lim_{h\to 0'_{\mathbb{X}}} f_{\mathbb{Y}}^{-1}\left(\frac{f_{\mathbb{Y}}\circ A\circ f_{\mathbb{X}}^{-1}\big[f_{\mathbb{X}}(x)+f_{\mathbb{X}}(h)\big] - f_{\mathbb{Y}}\circ A\circ f_{\mathbb{X}}^{-1}\big[f_{\mathbb{X}}(x)\big]}{f_{\mathbb{X}}(h)}\right) \tag{48}$$

$$= f_{\mathbb{Y}}^{-1}\left(\lim_{h\to 0}\frac{f_{\mathbb{Y}}\circ A\circ f_{\mathbb{X}}^{-1}\big[f_{\mathbb{X}}(x)+h\big] - f_{\mathbb{Y}}\circ A\circ f_{\mathbb{X}}^{-1}\big[f_{\mathbb{X}}(x)\big]}{h}\right) \tag{49}$$

$$= f_{\mathbb{Y}}^{-1}\left(\frac{d}{df_{\mathbb{X}}(x)}\underbrace{f_{\mathbb{Y}}\circ A\circ f_{\mathbb{X}}^{-1}}_{a}\big[f_{\mathbb{X}}(x)\big]\right) \tag{50}$$

$$= f_{\mathbb{Y}}^{-1}\left(\frac{da\big[f_{\mathbb{X}}(x)\big]}{df_{\mathbb{X}}(x)}\right). \tag{51}$$

This proves (22).

Now, let $A:\mathbb{X}\to\mathbb{Y}$, and define

$$\int_Y^X A(x)Dx = f_{\mathbb{Y}}^{-1}\left(\int_{f_{\mathbb{X}}(Y)}^{f_{\mathbb{X}}(X)} a(x)dx\right) \tag{52}$$

$$= f_{\mathbb{Y}}^{-1}\left(\int_{f_{\mathbb{X}}(Y)}^{f_{\mathbb{X}}(X)} f_{\mathbb{Y}}\circ A\circ f_{\mathbb{X}}^{-1}(x)dx\right). \tag{53}$$

Denoting $b(y)=\int_{f_{\mathbb{X}}(Y)}^{y} a(x)dx$ we rewrite

$$\int_Y^X A(x)Dx = f_{\mathbb{Y}}^{-1}\Big(b\big(f_{\mathbb{X}}(X)\big)\Big), \tag{54}$$

and thus (22) implies

$$\frac{D}{DX}\int_Y^X A(x)Dx = f_{\mathbb{Y}}^{-1}\left(\frac{db\big(f_{\mathbb{X}}(X)\big)}{df_{\mathbb{X}}(X)}\right) \tag{55}$$

$$= f_{\mathbb{Y}}^{-1}\Big(a\big(f_{\mathbb{X}}(X)\big)\Big) \tag{56}$$

$$= f_{\mathbb{Y}}^{-1}\circ a\circ f_{\mathbb{X}}(X) \tag{57}$$

$$= f_{\mathbb{Y}}^{-1}\circ f_{\mathbb{Y}}\circ A\circ f_{\mathbb{X}}^{-1}\circ f_{\mathbb{X}}(X) = A(X). \tag{58}$$

And the other way around,

$$\begin{aligned}
\int_Y^X \frac{DA(x)}{Dx} Dx &= \int_Y^X f_{\mathbb{Y}}^{-1} \circ a' \circ f_{\mathbb{X}}(x) Dx && (59)\\
&= f_{\mathbb{Y}}^{-1} \left(\int_{f_{\mathbb{X}}(Y)}^{f_{\mathbb{X}}(X)} f_{\mathbb{Y}} \circ f_{\mathbb{Y}}^{-1} \circ a' \circ f_{\mathbb{X}} \circ f_{\mathbb{X}}^{-1}(x) dx \right) && (60)\\
&= f_{\mathbb{Y}}^{-1} \left(\int_{f_{\mathbb{X}}(Y)}^{f_{\mathbb{X}}(X)} a'(x) dx \right) && (61)\\
&= f_{\mathbb{Y}}^{-1} \Big(a\big(f_{\mathbb{X}}(X)\big) - a\big(f_{\mathbb{X}}(Y)\big) \Big) && (62)\\
&= f_{\mathbb{Y}}^{-1} \Big(f_{\mathbb{Y}} \circ A \circ f_{\mathbb{X}}^{-1}\big(f_{\mathbb{X}}(X)\big) - f_{\mathbb{Y}} \circ A \circ f_{\mathbb{X}}^{-1}\big(f_{\mathbb{X}}(Y)\big) \Big) && (63)\\
&= f_{\mathbb{Y}}^{-1} \Big(f_{\mathbb{Y}}\big(A(X)\big) - f_{\mathbb{Y}}\big(A(Y)\big) \Big) && (64)\\
&= A(X) \ominus_{\mathbb{Y}} A(Y) && (65)
\end{aligned}$$